

Traits and alleles relevant for breeding and genetics:

last update May, 21st 2013

Associated markers, their chromosomal localisation, and the donor genotype/species are given. Chromosomal position of a trait/allele is given in megabases according to the 12 x genome sequence of PN40024 (<http://www.genoscope.cns.fr/vitis>). The symbols were discussed at the International Conference on Grapevine Breeding and Genetics at Geneva August 1 - 5, 2010 and assigned. Follow up information on naming of loci will be provided on VIVC to avoid homonyms.

For updating information please contact: reinhard.toepfer@jki.bund.de.

Symbol	Trait/Allele	Associated marker	Chromosome	Position on chromosome [Mb]	Parent 1	Parent 2	Population size	Genotype of origin	Original species trait/allele derived from?	Reference	Comment				
<i>Be size</i>	berry size (berry weight)	SCC8	18	25.9	MTP2223-27	x MTP2121-30	139		<i>V. vinifera</i>	Doligez et al. (2002)	Only one major QTL for berry size is indicated. There are several other QTLs described in the literature.				
		VMC7f2	18	26.9	Dominga	x Autumn Seedless	118			Cabezas et al. (2006)					
					Ruby Seedless	x Thompson Seedless	144			Mejia et al. (2007)					
<i>Mtc</i>	monoterpene content	DXS1	5	3.8	Italia	x Big Perlon	163			Costantini et al. (2008)					
						x Big Perlon	163		<i>V. vinifera</i>	Battilana et al. (2009)					
					Moscato Bianco	x <i>V. riparia</i>	174								
					Muscat Ottonel	x Muscat Ottonel	121		Duchene et al. (2009)						
<i>Lin</i>	Linalool content	cnd41	10		Italia	x Big Perlon	163		<i>V. vinifera</i>	Battilana et al. (2009)					
						x <i>V. riparia</i>	174								
						x Muscat Ottonel	121		Duchene et al. (2009)						
						x Gewürztraminer	115								
<i>Ffb</i>	Fleshless berry	VMC2A3	18	0.9	Chardonnay	x Ugni Blanc Mutant	71	Ugni Blanc	<i>V. vinifera</i>	Fernandez et al. (2006)	Mutant				
									<i>V. vinifera</i>						
<i>MybA</i>	berry skin colour	VMCNg3h8	14	26.6	V. rupestris	x V. arizonica	181		<i>V. arizonica</i>	Riaz et al. (2006)					
										VVIn64					Riaz et al. (2008)
										UDV-095		26.1			
<i>Rdv1</i>	<i>Daktulosphaira vitifoliae</i>	Gf13_9	13	21.9	Gf.V3125	x Börner	188	Börner	<i>V. cinerea</i>	Zhang et al. (2009)					
		VMC8e6		22.5											
<i>Rpv1</i>	<i>Plasmopara viticola</i>	VMC72	12		Syrah	x 28-8-78		28-8-78	<i>M. rotundifolia</i>	Merdinoglu et al. (2003)					
		VV1b32		10.3											
<i>Rpv2</i>	<i>Plasmopara viticola</i>		18		Cabernet Sauvignon	x 8624	129	8624	<i>M. rotundifolia</i>	Wiedemann-Merdinoglu et al. (2006)					
										Bellin et al. (2010)					
<i>Rpv3</i>	<i>Plasmopara viticola</i>	UDV-112	18		Regent	x Lemberger	153	Regent		Welter et al. (2007)					
										UDV-305		24.9			Bellin et al. (2009)
										VMC7f2		26.9			
<i>Rpv4</i>	<i>Plasmopara viticola</i>	VMC7h3	4	4.7	Regent	x Lemberger	153	Regent		Welter et al. (2007)					
		VMCNg2e1		5.2											
<i>Rpv5</i>	<i>Plasmopara viticola</i>	VV1o52b	9	4.0	Cabernet Sauvignon	x Gloire de Montpellier	138	Gloire de Montpellier	<i>V. riparia</i>	Marguerit et al. (2009)					
<i>Rpv6</i>	<i>Plasmopara viticola</i>	VMC8G9	12	20.4	Cabernet Sauvignon	x Gloire de Montpellier	138		<i>V. riparia</i>	Marguerit et al. (2009)					
<i>Rpv7</i>	<i>Plasmopara viticola</i>	UDV-097	7	11.4	Chardonnay	x Bianca	116	Bianca		Bellin et al. (2009)					
<i>Rpv8</i>	<i>Plasmopara viticola</i>	Chr14V015	14	6.6	<i>V. amurensis</i> 'Ruprecht'	x <i>V. amurensis</i> 'Ruprecht'	232	<i>V. amurensis</i> 'Ruprecht'	<i>V. amurensis</i>	Blasi et al. (2011)					
<i>Rpv9</i>	<i>Plasmopara viticola</i>	CCoAOMT	7	16.6	Moscato Bianco	x <i>V. riparia</i>	174	Wr63	<i>V. riparia</i>	Moreira et al. (2011)	CCoAOMT is the candidate gene from which the marker IN0006 was derived				
<i>Rpv10</i>	<i>Plasmopara viticola</i>	GF09-46	9	3.7	Gf.Ga-52-42	x Solaris	256	Solaris	<i>V. amurensis</i>	Schwander et al. (2012)					
<i>Rpv11</i>	<i>Plasmopara viticola</i>	VVMD27	5	4.5	Regent	x Lemberger	153	Regent		Fischer et al. (2004)					
		CS1E104J11F			Chardonnay	x Bianca	116	Chardonnay		Bellin et al. (2009)					
		VCHR05C		4.1	Gf.Ga-52-42	x Solaris	256	Solaris		Schwander et al. (2011)					
<i>Rpv12</i>	<i>Plasmopara viticola</i>	UDV-014	14	8.0	99-1-48	x Pinot noir	180	99-1-48	<i>V. amurensis</i>	Venuti et al. (2013)					
		UDV-304		9.3	Cabernet Sauvignan	x 20/3		20/3	<i>V. amurensis</i>						
		rgyvin180													
<i>Rpv13</i>	<i>Plasmopara viticola</i>	UDV-370		10.1											
		VMC1G3.2	12	10.0	Moscato Bianco	x <i>V. riparia</i>	174	Wr63	<i>V. riparia</i>	Moreira et al. (2011)					

Symbol	Trait/Allele	Associated marker	Chromosome	Position on chromosome [Mb]	Parent 1	Parent 2	Population size	Genotype of origin	Original species trait/allele derived from?	Reference	Comment
<i>Rcg1</i>	<i>Agrobacterium spec.</i>	UDV-015 9M3-3	15	7.1 9.3	Kunbarát	x Sárfehér	272	Kunbarát	<i>V. amurensis</i>	Kuczmozg et al. (2012)	
<i>Ren1</i>	<i>Erysiphe necator</i>	UDV-020 VMC9h4-2 VMCNg4e10.1	13	18.4 18.4	Nimrang	x Kishmish vatkana	310	Kishmish vatkana		Hoffmann et al. (2008)	
<i>Ren2</i>	<i>Erysiphe necator</i>	CS25	14	26.9	Horizon	x Illinois 547-1	58	Illinois 547-1		Dalbo et al. (2001)	
<i>Ren3</i>	<i>Erysiphe necator</i>	UDV-015b VViv67	15	7.1 10.9	Regent	x Lemberger	153	Regent		Welter et al. (2007)	
<i>Ren4</i>	<i>Erysiphe necator</i>	VMC7f2 SNPs	18	26.9 26.9	C166-043 C87-41	x F8909-08 x B70-57	42 57	C166-043 C87-41	<i>V. romanetii</i> <i>V. romanetii</i>	Riaz et al. (2012) Mahani et al. (2012)	
<i>Ren5</i>	<i>Erysiphe necator</i>		14	4.8					<i>M. rotundifolia</i>	Blanc et al. (2012)	
<i>Ren6</i>	<i>Erysiphe necator</i>									Barba et al. submitted	
<i>Run1</i>	<i>Erysiphe (Uncinula) necator</i>	VMC4f3.1 VMC8g9	12	13.1 20.4	VRH3082-1-42	x Cabernet Sauvignon	161	VRH3082-1-42	<i>M. rotundifolia</i>	Barker et al. (2005)	powdery mildew resistance originating from <i>Muscaninia</i> should be named as <i>Run...</i>
<i>Run2.1</i>	<i>Erysiphe (Uncinula) necator</i>	VMC7f2 VMCNg1e3 VVIn16 VMC7f2 VMC7f2	18	26.9 20.9 23.4 26.9 26.9	JB81-107-11	x Chenin Blanc x Tokay	97 47	Magnolia	<i>M. rotundifolia</i>	Riaz et al. (2011)	resistant tissue: Cane Rachis Rachis Fruit Leaf, Cane, Rachis, Fruit
<i>Run2.2</i>	<i>Erysiphe (Uncinula) necator</i>	VMC7f2	18	26.9	A90-71 e2-9	x Flame Seedless x Malaga Rosada	80 255	Trayshed	<i>M. rotundifolia</i>	Riaz et al. (2011)	
<i>Sd1</i>	seed development inhibitor	SCC8 VMC6f11	18	25.9 23.2	MTP2223-27 Dominga	x MTP2121-30 x Autumn Seedless	139 118	Autumn Seedless		Doligez et al. (2002) Cabezas et al. (2006)	
	seedlessness	VMC7f2		26.9	Dominga Italia	x Autumn Seedless x Big Perlon	118 163	Big Perlon		Costantini et al. (2008) Barba et al. submitted	
<i>Sen1</i>											
<i>Sex</i>	sex	VVMD34 VVS3 VVib23	2	3.7 4.2 4.9	Horizon Ramsey	x Illinois 547-1 x Riparia Gloire	58 188			Dalbó et al. (2000) Lowe and Walker (2006)	
				5.0	<i>V. rupestris</i> V3125	x <i>V. arizonica</i> x Börner	181 202			Riaz et al. (2006) Fechter et al. (2012)	
<i>Ufgt</i>		UFGT	16	2.3	Regent	x Lemberger	153			Fischer et al. (2004)	
<i>Ver</i>	véraison	VMC1E11	16	13.7	Regent Italia	x Lemberger x Big Perlon	153 163	Regent		Fischer et al. (2004) Costantini et al. (2008)	For véraison (begin of ripening) several QTLs are published. The locus on LG 16 is the only one which was found in two independent mapping populations.
<i>Vygail</i>	GA insensitive dwarf mutant		1	4.9				Pinot Meunier		Boss & Thomas (2002)	
<i>Xir1</i>	<i>Xiphinema index</i>	VMC5a10 1N2R3b M4F3R	19	20.9 20.9	<i>V. rupestris</i> <i>V. rupestris</i>	x <i>V. arizonica</i> x <i>V. arizonica</i>	185 185		<i>V. arizonica</i> <i>V. arizonica</i>	Xu et al. (2008) Hwang et al. (2010)	
<i>5-gt</i>	anthocyanin 3,5-diglucosides	Gf09_01	9	6.5	Regent	x Lemberger	153	Regent		Hausmann et al. (2009)	